arXiv:math/0501038v4 [math.GM] 11 Jan 2006

THE MASLOV DEQUANTIZATION, IDEMPOTENT AND TROPICAL MATHEMATICS: A VERY BRIEF INTRODUCTION

G. L. LITVINOV

Abstract. This paper is a very brief introduction to idempotent mathematics and related topics. It appears as an introductory paper in the volume *Idempotent Mathematics and Mathematical Physics* (G. L. Litvinov and V. P. Maslov, eds; AMS Contemporary Mathematics Proceedings Series, 2005) [73].

This paper is a very brief introduction, without exact theorems and proofs, to the Maslov dequantization and idempotent and tropical mathematics. Our list of references is not complete (not at all). Additional references can be found, e.g., in the electronic archive

`http://arXiv.org`,

in [6, 10, 13, 16, 19–22, 26, 27, 35, 42, 44–49, 56, 57, 62, 65, 70, 72, 78, 83, 137] and in the papers published in this volume.

The author thanks M. Akian, Ya. I. Belopolskaya, P. Butkovič, G. Cohen, S. Gaubert, R. A. Guninghame-Green, P. Del Moral, H. Prade, W. H. Fleming, J. S. Golan, M. Gondran, I. Itenberg, Y. Katsov, V. N. Kolokoltsov, P. Loreti, Yu. I. Manin, G. Mascari, W. M. MacEneaney, E. Pap, M. Pedicini, A. A. Puhalskii, J.-P. Quadrat, M. A. Roytberg, G. B. Shpiz, I. Singer, and O. Viro for useful contacts and references. Special thanks are due to V. P. Maslov for his crucial help and support and to A. N. Sobolevskiĭ for his great help including, but not limited to, two pictures presented below.

2000 *Mathematics Subject Classification.* Primary 00A05, 81Q20, 14P99, 51P05, 46S10, 49L99; Secondary 06A99, 06F07, 14N10, 81S99.

Key words and phrases. The Maslov dequantization, idempotent mathematics, tropical mathematics, idempotent semirings, idempotent analysis, idempotent functional analysis, dequantization of geometry.

This work has been supported by the RFBR grant 05–01–00824 and the Erwin Schrödinger International Institute for Mathematical Physics (ESI).

1. Some basic ideas

Idempotent mathematics is based on replacing the usual arithmetic operations with a new set of basic operations (such as maximum or minimum), that is on replacing numerical fields by idempotent semirings and semifields. Typical examples are given by the so-called max-plus algebra $\mathbf{R}_{\max}$ and the min-plus algebra $\mathbf{R}_{\min}$. Let $\mathbf{R}$ be the field of real numbers. Then $\mathbf{R}_{\max} = \mathbf{R} \cup \{-\infty\}$ with operations $x \oplus y = \max\{x, y\}$ and $x \odot y = x + y$. Similarly $\mathbf{R}_{\min} = \mathbf{R} \cup \{+\infty\}$ with the operations $\oplus = \min$, $\odot = +$. The new addition $\oplus$ is idempotent, i.e., $x \oplus x = x$ for all elements x.

Many authors (S. C. Kleene, N. N. Vorobjev, B. A. Carre, R. A. Cuninghame-Green, K. Zimmermann, U. Zimmermann, M. Gondran, F. L. Baccelli, G. Cohen, S. Gaubert, G. J. Olsder, J.-P. Quadrat, and others) used idempotent semirings and matrices over these semirings for solving some applied problems in computer science and discrete mathematics, starting from the classical paper of S. C. Kleene [58]. The modern *idempotent analysis* (or *idempotent calculus*, or *idempotent mathematics*) was founded by V. P. Maslov in the 1980s in Moscow; see, e.g., [65, 87–92].

Idempotent mathematics can be treated as a result of a dequantization of the traditional mathematics over numerical fields as the Planck constant $\hbar$ tends to zero taking imaginary values. This point of view was presented by G. L. Litvinov and V. P. Maslov [70–72], see also [78, 79]. In other words, idempotent mathematics is an asymptotic version of the traditional mathematics over the fields of real and complex numbers.

The basic paradigm is expressed in terms of an *idempotent correspondence principle.* This principle is closely related to the well-known correspondence principle of N. Bohr in quantum theory. Actually, there exists a heuristic correspondence between important, interesting, and useful constructions and results of the traditional mathematics over fields and analogous constructions and results over idempotent semirings and semifields (i.e., semirings and semifields with idempotent addition).

A systematic and consistent application of the idempotent correspondence principle leads to a variety of results, often quite unexpected. As a result, in parallel with the traditional mathematics over fields, its “shadow,” the idempotent mathematics, appears. This “shadow” stands approximately in the same relation to the traditional mathematics as does classical physics to quantum theory, see Fig. 1. In many respects idempotent mathematics is simpler than the traditional one.


TRADITIONAL MATHEMATICS
Fields of real and complex numbers
Idempotent Correspondence Principle
IDEMPOTENT MATHEMATICS
Idempotent semirings and semifields
N. Bohr's Correspondence Principle
QUANTUM MECHANICS
CLASSICAL MECHANICS

Figure 1

However the transition from traditional concepts and results to their idempotent analogs is often nontrivial.

2. Semirings, semifields, and dequantization

Consider a set S equipped with two algebraic operations: *addition* $\oplus$ and *multiplication* $\odot$. It is a *semiring* if the following conditions are satisfied:

- the addition $\oplus$ and the multiplication $\odot$ are associative;
- the addition $\oplus$ is commutative;
- the multiplication $\odot$ is distributive with respect to the addition $\oplus$:

$$x \odot (y \oplus z) = (x \odot y) \oplus (x \odot z) \quad \text{and} \quad (x \oplus y) \odot z = (x \odot z) \oplus (y \odot z)$$

for all $x, y, z \in S$.

A *unity* of a semiring S is an element $\mathbf{1} \in S$ such that $\mathbf{1} \odot x = x \odot \mathbf{1} = x$ for all $x \in S$. A *zero* of a semiring S is an element $\mathbf{0} \in S$ such that $\mathbf{0} \neq \mathbf{1}$ and $\mathbf{0} \oplus x = x$, $\mathbf{0} \odot x = x \odot \mathbf{0} = \mathbf{0}$ for all $x \in S$. A semiring S is called an *idempotent semiring* if $x \oplus x = x$ for all $x \in S$. A semiring S with neutral elements $\mathbf{0}$ and $\mathbf{1}$ is called a *semifield* if every nonzero element of S is invertible. Note that dioïds in the sense of [6, 46, 47], quantales in the sense of [109, 110], and inclines in the sense of [56] are examples of idempotent semirings.

Let $\mathbf{R}$ be the field of real numbers and $\mathbf{R}_+$ the semiring of all nonnegative real numbers (with respect to the usual addition and multiplication). The change of variables $x \mapsto u = h \ln x$, $h > 0$, defines a map $\Phi_h \colon \mathbf{R}_+ \to S = \mathbf{R} \cup \{-\infty\}$. Let the addition and multiplication operations be mapped from $\mathbf{R}$ to S by Φ_h, i.e., let $u \oplus_h v = h \ln(\exp(u/h) + \exp(v/h))$, $u \odot v = u + v$, $\mathbf{0} = -\infty = \Phi_h(0)$, $\mathbf{1} = 0 = \Phi_h(1)$. It can easily be checked that $u \oplus_h v \to \max\{u, v\}$ as $h \to 0$ and that S forms a semiring with respect to addition $u \oplus v = \max\{u, v\}$ and multiplication $u \odot v = u + v$ with zero $\mathbf{0} = -\infty$ and unit $\mathbf{1} = 0$. Denote this semiring by $\mathbf{R}_{\max}$; it is *idempotent*, i.e., $u \oplus u = u$ for all its elements. The semiring $\mathbf{R}_{\max}$ is actually a semifield. The analogy with quantization is obvious; the parameter h plays the rôle of the Planck constant, so $\mathbf{R}_+$ (or $\mathbf{R}$) can be viewed as a "quantum object" and $\mathbf{R}_{\max}$ as the result of its "dequantization." A similar procedure (for $h < 0$) gives the semiring $\mathbf{R}_{\min} = \mathbf{R} \cup \{+\infty\}$ with the operations $\oplus = \min$, $\odot = +$; in this case $\mathbf{0} = +\infty$, $\mathbf{1} = 0$. The semirings $\mathbf{R}_{\max}$ and $\mathbf{R}_{\min}$ are isomorphic. This passage to $\mathbf{R}_{\max}$ or $\mathbf{R}_{\min}$ is called the *Maslov dequantization.* It is clear that the corresponding passage from $\mathbf{C}$ or $\mathbf{R}$ to $\mathbf{R}_{\max}$ is generated by the Maslov dequantization and the map $x \mapsto |x|$. By misuse of language, *we shall also call this passage the Maslov dequantization.* Connections with physics and the meaning of imaginary values of the Planck constant are discussed in [78, 79]. The idempotent semiring $\mathbf{R} \cup \{-\infty\} \cup \{+\infty\}$ with the operations $\oplus = \max$, $\odot = \min$ can be obtained as a result of a "second dequantization" of $\mathbf{C}$, $\mathbf{R}$ or $\mathbf{R}_+$. Dozens of interesting examples of nonisomorphic idempotent semirings may be cited as well as a number of standard methods of deriving new semirings from these (see, e.g., [17, 44–49, 72, 78]). The so-called *idempotent dequantization* is a generalization of the Maslov dequantization; this is a passage from fields to idempotent semifields and semirings in mathematical constructions and results.

The Maslov dequantization is related to the well-known logarithmic transformation that was used, e.g., in the classical papers of E. Schrödinger (1926) and E. Hopf (1951). The term 'Cole-Hopf transformation' is also used. The subsequent progress of E. Hopf's ideas has culminated in the well-known vanishing viscosity method and the method of viscosity solutions, see, e.g., [7, 12, 38, 90, 122] and papers [93] by D. McCaffrey and [111] by I. V. Roublev published in this volume.

3. TERMINOLOGY: TROPICAL SEMIRINGS AND TROPICAL MATHEMATICS

The term 'tropical semirings' was introduced in computer science to denote discrete versions of the max-plus algebra $\mathbf{R}_{\max}$ or min-plus algebra $\mathbf{R}_{\min}$ and their subalgebras; (discrete) semirings of this type were called tropical semirings by Dominic Perrin in honour of Imre Simon (who is a Brasilian mathematician and computer scientist) because of his pioneering activity in this area, see [102].

More recently the situation and terminology have changed. For the most part of modern authors ‘tropical’ means ‘over $\mathbf{R}_{\max}$ (or $\mathbf{R}_{\min}$)’ and tropical semirings are idempotent semifields $\mathbf{R}_{\max}$ and $\mathbf{R}_{\min}$. The terms ‘max-plus’ and ‘min-plus’ are often used in the same sense. Now the term ‘tropical mathematics’ usually means ‘mathematics over $\mathbf{R}_{\max}$ or $\mathbf{R}_{\min}$’, see, e.g., [50, 94–97, 108, 120]. Terms ‘tropicalization’ and ‘tropification’ (see, e.g., [57]) mean exactly dequantization and quantization in our sense. In any case, tropical mathematics is a natural and very important part of idempotent mathematics.

Note that in papers [125–127] N. N. Vorobjev developed a version of idempotent linear algebra (with important applications, e.g., to mathematical economics) and predicted many aspects of the future extended theory. He used the terms ‘extremal algebras’ and ‘extremal mathematics’ for idempotent semirings and idempotent mathematics. Unfortunately, N. N. Vorobjev’s papers and ideas were forgotten for a long period, so his remarkable terminology is not in use any more.

4. Idempotent algebra and linear algebra

The first known paper on idempotent (linear) algebra is due to S. Kleene [58]. Systems of linear algebraic equations over an exotic idempotent semiring of all formal languages over a fixed finite alphabet are examined in this work; however, S. Kleene’s ideas are very general and universal. Since then, dozens of authors investigated matrices with coefficients belonging to an idempotent semiring and the corresponding applications to discrete mathematics, computer science, computer languages, linguistic problems, finite automata, optimization problems on graphs, discrete event systems and Petry nets, stochastic systems, computer performance evaluation, computational problems etc. This subject is very well known and well presented in the corresponding literature, see, e.g., [6, 10, 11, 13, 16, 19–21, 31, 42, 44–49, 56, 62, 65, 70, 72–75, 82, 83, 90, 125–127, 137]. The idempotent linear algebra is treated in the papers of P. Butkovič [11] and E. Wagneur [128] in the present volume.

Idempotent abstract algebra is not so well developed yet (on the other hand, from a formal point of view, the lattice theory and the theory of ordered groups and semigroups are parts of idempotent algebra). However, there are many interesting results and applications presented, e.g., in [20–22, 53, 109, 110, 114].

In particular, an idempotent version of the main theorem of algebra holds [22, 114] for radicable idempotent semifields (a semiring A is *radicable* if the equation $x^n = a$ has a solution $x \in A$ for any $a \in A$

and any positive integer n). It is proved that $\mathbf{R}_{\max}$ and other radicable semifields are algebraically closed in a natural sense [114].

5. Idempotent analysis

Idempotent analysis was initially constructed by V. P. Maslov and his collaborators and then developed by many authors. The subject is presented in the book of V. N. Kolokoltsov and V. P. Maslov [65] (a version of this book in Russian [90] was published in 1994).

Let S be an arbitrary semiring with idempotent addition $\oplus$ (which is always assumed to be commutative), multiplication $\odot$, zero $\mathbf{0}$, and unit $\mathbf{1}$. The set S is supplied with the *standard partial order* $\preceq$: by definition, $a \preceq b$ if and only if $a \oplus b = b$. Thus all elements of S are nonnegative: $\mathbf{0} \preceq a$ for all $a \in S$. Due to the existence of this order, idempotent analysis is closely related to the lattice theory, theory of vector lattices, and theory of ordered spaces. Moreover, this partial order allows to model a number of basic "topological" concepts and results of idempotent analysis at the purely algebraic level; this line of reasoning was examined systematically in [76–80] and [17].

Calculus deals mainly with functions whose values are numbers. The idempotent analog of a numerical function is a map $X \to S$, where X is an arbitrary set and S is an idempotent semiring. Functions with values in S can be added, multiplied by each other, and multiplied by elements of S pointwise.

The idempotent analog of a linear functional space is a set of S-valued functions that is closed under addition of functions and multiplication of functions by elements of S, or an S-semimodule. Consider, e.g., the S-semimodule $B(X, S)$ of all functions $X \to S$ that are bounded in the sense of the standard order on S.

If $S = \mathbf{R}_{\max}$, then the idempotent analog of integration is defined by the formula

$$I(\varphi) = \int_X^{\oplus} \varphi(x)\, dx = \sup_{x \in X} \varphi(x), \tag{1}$$

where $\varphi \in B(X, S)$. Indeed, a Riemann sum of the form $\sum\limits_i \varphi(x_i) \cdot \sigma_i$ corresponds to the expression $\bigoplus\limits_i \varphi(x_i) \odot \sigma_i = \max\limits_i \{\varphi(x_i) + \sigma_i\}$, which tends to the right-hand side of (1) as $\sigma_i \to 0$. Of course, this is a purely heuristic argument.

Formula (1) defines the *idempotent* (or *Maslov*) *integral* not only for functions taking values in $\mathbf{R}_{\max}$, but also in the general case when any of bounded (from above) subsets of S has the least upper bound.

An *idempotent* (or *Maslov*) *measure* on X is defined by $m_\psi(Y) = \sup_{x \in Y} \psi(x)$, where $\psi \in B(X,S)$. The integral with respect to this measure is defined by

$$I_\psi(\varphi) = \int_X^\oplus \varphi(x)\, dm_\psi = \int_X^\oplus \varphi(x) \odot \psi(x)\, dx = \sup_{x \in X} (\varphi(x) \odot \psi(x)). \quad (2)$$

Obviously, if $S = \mathbf{R}_{\min}$, then the standard order $\preceq$ is opposite to the conventional order $\leq$, so in this case equation (2) assumes the form

$$\int_X^\oplus \varphi(x)\, dm_\psi = \int_X^\oplus \varphi(x) \odot \psi(x)\, dx = \inf_{x \in X} (\varphi(x) \odot \psi(x)),$$

where inf is understood in the sense of the conventional order $\leq$.

Note that the so-called pseudo-analysis (see a survey paper of E. Pap [100] published in the present volume) is related to a special part of idempotent analysis; however, this pseudo-analysis is not a proper part of idempotent mathematics in the general case.

6. CORRESPONDENCE TO STOCHASTICS AND A DUALITY BETWEEN PROBABILITY AND OPTIMIZATION

Maslov measures are nonnegative (in the sense of the standard order) just as probability measures. The analogy between idempotent and probability measures leads to important relations between optimization theory and probability theory. By the present time idempotent analogues of many objects of stochastic calculus have been constructed and investigated, such as max-plus martingales, max-plus stochastic differential equations, and others. These results allow, for example, to transfer powerful stochastic methods to the optimization theory. This was noticed and examined by many authors (G. Salut, P. Del Moral, M. Akian, J.-P. Quadrat, V. P. Maslov, V. N. Kolokoltsov, P. Bernhard, W. A. Fleming, W. M. McEneaney, A. A. Puhalskii and others), see the survey paper of W. A. Fleming and W. M. McEneaney [37] published in this volume and [1, 4, 8, 19, 24–27, 34–36, 48, 90, 103, 105, 106]. For relations and applications to large deviations see [1, 24–27, 104] and especially the book of A. A. Puhalskii [103].

7. IDEMPOTENT FUNCTIONAL ANALYSIS

Many other idempotent analogs may be given, in particular, for basic constructions and theorems of functional analysis. Idempotent functional analysis is an abstract version of idempotent analysis. For the sake of simplicity take $S = \mathbf{R}_{\max}$ and let X be an arbitrary set. The idempotent integration can be defined by the formula (1), see above.

The functional $I(\varphi)$ is linear over S and its values correspond to limiting values of the corresponding analogs of Lebesgue (or Riemann) sums. An idempotent scalar product of functions φ and ψ is defined by the formula

$$\langle \varphi, \psi \rangle = \int_X^{\oplus} \varphi(x) \odot \psi(x)\, dx = \sup_{x \in X} (\varphi(x) \odot \psi(x)).$$

So it is natural to construct idempotent analogs of integral operators in the form

$$K : \varphi(y) \mapsto (K\varphi)(x) = \int_Y^{\oplus} K(x,y) \odot \varphi(y)\, dy = \sup_{y \in Y} \{K(x,y) + \varphi(y)\}, \tag{3}$$

where $\varphi(y)$ is an element of a space of functions defined on a set Y, and $K(x,y)$ is an S-valued function on $X \times Y$. Of course, expressions of this type are standard in optimization problems.

Recall that the definitions and constructions described above can be extended to the case of idempotent semirings which are conditionally complete in the sense of the standard order. Using the Maslov integration, one can construct various function spaces as well as idempotent versions of the theory of generalized functions (distributions). For some concrete idempotent function spaces it was proved that every 'good' linear operator (in the idempotent sense) can be presented in the form (3); this is an idempotent version of the kernel theorem of L. Schwartz; results of this type were proved by V. N. Kolokoltsov, P. S. Dudnikov and S. N. Samborski, I. Singer, M. A. Shubin and others, see, e.g., [31, 65, 90, 91, 117]. So every 'good' linear functional can be presented in the form $\varphi \mapsto \langle \varphi, \psi \rangle$, where $\langle , \rangle$ is an idempotent scalar product.

In the framework of idempotent functional analysis results of this type can be proved in a very general situation. In [76–80] an algebraic version of the idempotent functional analysis is developed; this means that basic (topological) notions and results are simulated in purely algebraic terms. The treatment covers the subject from basic concepts and results (e.g., idempotent analogs of the well-known theorems of Hahn-Banach, Riesz, and Riesz-Fisher) to idempotent analogs of A. Grothendieck's concepts and results on topological tensor products, nuclear spaces and operators. An abstract version of the kernel theorem is formulated. Note that the passage from the usual theory to idempotent functional analysis may be very nontrivial; for example, there are many non-isomorphic idempotent Hilbert spaces. Important results on idempotent functional analysis (duality and separation theorems) are recently published by G. Cohen, S. Gaubert, and J.-P. Quadrat [17]; see also a finite dimensional version of the separation theorem in [133]. Three papers on this subject by M. Akian, S. Gaubert, V. Kolokoltsov, G. Cohen, J.-P. Quadrat, I. Singer, and C. Walsh [3, 5, 18] are published in this volume.

There is an "idempotent" version of the theory of linear representations of groups and semigroups and the abstract harmonic analysis, see, e.g., [79]. In the framework of this theory the well-known Legendre transform can be treated as an idempotent version of the traditional Fourier transform (this observation is due to V. P. Maslov).

8. The superposition principle and linear problems

Basic equations of quantum theory are linear; this is the superposition principle in quantum mechanics. The Hamilton–Jacobi equation, the basic equation of classical mechanics, is nonlinear in the conventional sense. However, it is linear over the semirings $\mathbf{R}_{\max}$ and $\mathbf{R}_{\min}$. Similarly, different versions of the Bellman equation, the basic equation of optimization theory, are linear over suitable idempotent semirings; this is V. P. Maslov's idempotent superposition principle, see [87–91]. For instance, the finite-dimensional stationary Bellman equation can be written in the form $X = H \odot X \oplus F$, where X, H, F are matrices with coefficients in an idempotent semiring S and the unknown matrix X is determined by H and F. In particular, standard problems of dynamic programming and the well-known shortest path problem correspond to the cases $S = \mathbf{R}_{\max}$ and $S = \mathbf{R}_{\min}$, respectively. It is known that principal optimization algorithms for finite graphs correspond to standard methods for solving systems of linear equations of this type (i.e., over semirings). Specifically, Bellman's shortest path algorithm corresponds to a version of Jacobi's algorithm, Ford's algorithm corresponds to the Gauss–Seidel iterative scheme, etc.

The linearity of the Hamilton–Jacobi equation over $\mathbf{R}_{\min}$ and $\mathbf{R}_{\max}$, which is the result of the Maslov dequantization of the Schrödinger equation, is closely related to the (conventional) linearity of the Schrödinger equation and can be deduced from this linearity. Thus, it is possible to borrow standard ideas and methods of linear analysis and apply them to a new area.

The action functional $S = S(x(t))$ can be considered as a function taking the set of curves (trajectories) to the set of real numbers which can be treated as elements of $\mathbf{R}_{\min}$. In this case the minimum of the action functional can be viewed as the Maslov integral of this function over the set of trajectories or an idempotent analog of the Euclidean version of the Feynman path integral. The minimum of the action functional corresponds to the maximum of e^{-S}, i.e. idempotent integral $\int^{\oplus}_{\{paths\}} e^{-S(x(t))} D\{x(t)\}$. Thus the least action principle can be considered as an idempotent version of the well-known Feynman approach to quantum mechanics. The representation of a solution to the Schrödinger equation in terms of the Feynman integral corresponds to the Lax–Oleĭnik solution formula for the Hamilton–Jacobi equation.

The idempotent superposition principle indicates that there exist important nonlinear (in the traditional sense) problems that are linear over idempotent semirings. The linear idempotent functional analysis is a natural tool for investigation of those nonlinear infinite-dimensional problems that possess this property.

9. Dequantization of geometry

An idempotent version of real algebraic geometry was discovered in the report of O. Viro for the Barcelona Congress [123]. Starting from the idempotent correspondence principle O. Viro constructed a piecewise-linear geometry of polyhedra of a special kind in finite dimensional Euclidean spaces as a result of the Maslov dequantization of real algebraic geometry. He indicated important applications in real algebraic geometry (e.g., in the framework of Hilbert's 16th problem for constructing real algebraic varieties with prescribed properties and parameters) and relations to complex algebraic geometry and amoebas in the sense of I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky (see their book [41] and [124]). Then complex algebraic geometry was dequantized by G. Mikhalkin and the result turned out to be the same; this new 'idempotent' (or asymptotic) geometry is now often called the *tropical algebraic geometry*, see, e.g., [32, 50, 94–97, 108, 115, 120, 121].

There is a natural relation between the Maslov dequantization and amoebas. Suppose $(\mathbf{C}^*)^n$ is a complex torus, where $\mathbf{C}^* = \mathbf{C}\backslash\{0\}$ is the group of nonzero complex numbers under multiplication. For $z = (z_1, \dots, z_n) \in (\mathbf{C}^*)^n$ and a positive real number h denote by $\mathrm{Log}_h(z) = h\log(|z|)$ the element

$$(h \log|z_1|, h\log|z_2|, \dots, h\log|z_n|) \in \mathbf{R}^n.$$

Suppose $V \subset (\mathbf{C}^*)^n$ is a complex algebraic variety; denote by $\mathcal{A}_h(V)$ the set $\mathrm{Log}_h(V)$. If $h = 1$, then the set $\mathcal{A}(V) = \mathcal{A}_1(V)$ is called the *amoeba* of V in the sense of [41], see also [124]; the amoeba $\mathcal{A}(V)$ is a closed subset of $\mathbf{R}^n$ with a non-empty complement. Note that this construction depends on our coordinate system.



FIGURE 2

For the sake of simplicity suppose V is a hypersurface in $(\mathbf{C}^*)^n$ defined by a polynomial f; then there is a deformation $h \mapsto f_h$ of this polynomial generated by the Maslov dequantization and $f_h = f$ for $h = 1$. Let $V_h \subset (\mathbf{C}^*)^n$ be the zero set of f_h and set $\mathcal{A}_h(V_h) = \mathrm{Log}_h(V_h)$. Then there exists a tropical variety $\mathit{Tro}(V)$ such that the subsets $\mathcal{A}_h(V_h) \subset \mathbf{R}^n$ tend to $\mathit{Tro}(V)$ in the Hausdorff metric as $h \to 0$, see [94, 113]. The tropical variety $\mathit{Tro}(V)$ is a result of a deformation of the amoeba $\mathcal{A}(V)$ and the Maslov dequantization of the variety V. The set $\mathit{Tro}(V)$ is called the *skeleton* of $\mathcal{A}(V)$.

Example [94]**.** For the line $V = \{\, (x, y) \in (\mathbf{C}^*)^2 \mid x + y + 1 = 0 \,\}$ the piecewise-linear graph $\mathit{Tro}(V)$ is a tropical line, see Fig. 2(a). The amoeba $\mathcal{A}(V)$ is represented in Fig. 2(b), while Fig. 2(c) demonstrates the corresponding deformation of the amoeba.

There is an interesting paper [101] of M. Passare and A. Tsikh on amoebas of algebraic and analytic varieties in the present volume.

In the important paper [52] (see also [32, 94, 96, 108]) tropical varieties appeared as amoebas over non-Archimedean fields. In 2000 M. Kontsevich noted that it is possible to use non-Arhimedean amoebas in enumerative geometry, see [94, section 2.4, remark 4]. In fact methods of tropical geometry lead to remarkable applications to the algebraic enumerative geometry, Gromov-Witten and Welschinger invariants, see [50, 51, 94–97]. In particular, G. Mikhalkin presented and proved in [95, 97] a formula enumerating curves of arbitrary genus in toric surfaces.

Note that tropical geometry is closely related to the well-known program of M. Kontsevich and Y. Soibelman, see, e.g., [67, 68]. There is an introductory paper [108] on tropical algebraic geometry in this volume. The paper of G. L. Litvinov and G. B. Shpiz [81] (which is also published in the present volume) is also related to the subject.

However on the whole only first steps in idempotent/tropical geometry have been made and the problem of systematic construction of idempotent versions of algebraic and analytic geometries is still open.

10. The correspondence principle for algorithms and their computer implementations

There are many important applied algorithms of idempotent mathematics, see, e.g., [6, 11, 13, 21, 22, 33, 36, 37, 46–48, 50, 56, 57, 62, 65, 70, 72, 74, 75, 82–84, 95, 97, 108, 112, 121, 125–127, 129, 136, 137]. The idempotent correspondence principle is valid for algorithms as well as for their software and hardware implementations [70–72, 74, 75]. In particular, according to the superposition principle, analogs of linear algebra algorithms are especially important. It is well-known that algorithms of linear algebra are convenient for parallel computations; so their idempotent analogs accept a parallelization. This is a regular way to use parallel computations for many problems including basic optimization problems. It is convenient to use universal algorithms which do not depend on a concrete semiring and its concrete computer model. Software implementations for universal semiring algorithms are based on object-oriented and generic programming; program modules can deal with abstract (and variable) operations and data types, see [70, 72, 74, 75]. The paper [84] of P. Loreti and M. Pedicini on the subject is published in the present volume.

The most important and standard algorithms have many hardware implementations in the form of technical devices or special processors. These devices often can be used as prototypes for new hardware units generated by substitution of the usual arithmetic operations for its semiring analogs, see [70, 72, 75]. Good and efficient technical ideas and decisions can be transposed from prototypes into new hardware units. Thus the correspondence principle generates a regular heuristic method for hardware design.

11. Idempotent interval analysis

An idempotent version of the traditional interval analysis is presented in [82, 83]. Let S be an idempotent semiring equipped with the standard partial order. A *closed interval* in S is a subset of the form $\mathbf{x} = [\underline{\mathbf{x}}, \bar{\mathbf{x}}] = \{x \in S \mid \underline{\mathbf{x}} \preceq x \preceq \bar{\mathbf{x}}\}$, where the elements $\underline{\mathbf{x}} \preceq \bar{\mathbf{x}}$ are called *lower* and *upper bounds* of the interval $\mathbf{x}$. A *weak interval extension* $I(S)$ of the semiring S is the set of all closed intervals in S endowed with operations $\oplus$ and $\odot$ defined as $\mathbf{x} \oplus \mathbf{y} = [\underline{\mathbf{x}} \oplus \underline{\mathbf{y}}, \bar{\mathbf{x}} \oplus \bar{\mathbf{y}}]$, $\mathbf{x} \odot \mathbf{y} = [\underline{\mathbf{x}} \odot \underline{\mathbf{y}}, \bar{\mathbf{x}} \odot \bar{\mathbf{y}}]$; the set $I(S)$ is a new idempotent semiring with respect to these operations. It is proved that basic interval problems of idempotent linear algebra are polynomial, whereas in the traditional interval analysis problems of this kind are generally NP-hard. Exact interval solutions for the discrete stationary Bellman equation (see the matrix equation discussed in section 8 above) and for the corresponding optimization problems are constructed and examined by G. L. Litvinov and A. Sobolevskiĭ in [82, 83]. Similar results are presented by K. Cechlárová and R. A. Cuninghame-Green in [14].

12. Relations to the KAM theory and optimal transport

The subject of the Kolmogorov–Arnold–Moser (KAM) theory may be formulated as the study of invariant subsets in phase spaces of nonintegrable Hamiltonian dynamical systems where the dynamics displays the same degree of regularity as that of integrable systems (quasiperiodic behaviour). Recently, a considerable progress was made via a variational approach, where the dynamics is specified by the Lagrangian rather than Hamiltonian function. The corresponding theory was initiated by S. Aubry and J. N. Mather and recently dubbed *weak KAM theory* by A. Fathi (see his book "Weak KAM Theorems in Lagrangian Dynamics," Cambridge Univ. Press, 2004; see also [54, 55, 118, 119]). Minimization of a certain functional along trajectories of moving particles is a central feature of another subject, the optimal transport theory, which also has undergone a rapid recent development. This theory dates back to G. Monge's work on cuts and fills (1781). A modern version of the theory is known now as the so-called *Monge–Ampère–Kantorovich (MAK) optimal transport theory* (after the work of L. V. Kantorovich "On the translocation of masses" in C.R. (Doklady) Acad. Sci. USSR, v. 321, 1942, p. 199–201). There is a similarity between the two theories and there are relations to problems of the idempotent functional analysis (e.g., the problem of eigenfunctions for "idempotent" integral operators, see [118]). Applications of optimal transport to data processing in cosmology are presented in [9, 40].

13. Relations to logic, fuzzy sets, and possibility theory

Let S be an idempotent semiring with neutral elements $\mathbf{0}$ and $\mathbf{1}$ (recall that $\mathbf{0} \neq \mathbf{1}$, see section 2 above). Then the Boolean algebra $\mathbf{B} = \{\mathbf{0}, \mathbf{1}\}$ is a natural idempotent subsemiring of S. Thus S can be treated as a generalized (extended) logic with logical operations $\oplus$ (disjuction) and $\odot$ (conjunction). Ideas of this kind are discussed in many books and papers with respect to generalized versions of logic and especially quantum logic, see, e.g., [44, 59, 109, 110]. In the present volume there is a paper of A. Di Nola and B. Gerla [29] related to these ideas.

Let Ω be the so-called universe consisting of "elementary events." Denote by $\mathcal{F}(S)$ the set of functions defined on Ω and taking their values in S; then $\mathcal{F}(S)$ is an idempotent semiring with respect to the pointwise addition and multiplication of functions. We shall say that elements of $\mathcal{F}(S)$ are *generalized fuzzy sets*, see [44, 69]. We have the well-known classical definition of fuzzy sets (L. A. Zadeh [130]) if $S = \mathbf{P}$, where $\mathbf{P}$ is the segment $[0, 1]$ with the semiring operations $\oplus = \max$ and $\odot = \min$. Of course, functions from $\mathcal{F}(\mathbf{P})$ taking their values in the Boolean algebra $\mathbf{B} = \{0, 1\} \subset \mathbf{P}$ correspond to traditional sets from Ω and semiring operations correspond to standard operations for sets. In the general case functions from $\mathcal{F}(S)$ taking their values in $\mathbf{B} = \{\mathbf{0}, \mathbf{1}\} \subset S$ can be treated as traditional subsets in Ω. If S is a lattice (i.e. $x \odot y = \inf\{x, y\}$ and $x \oplus y = \sup\{x, y\}$), then generalized fuzzy sets coincide with L-fuzzy sets in the sense of J. A. Goguen [43]. The set $I(S)$ of intervals is an idempotent semiring (see section 11), so elements of $\mathcal{F}(I(S))$ can be treated as interval (generalized) fuzzy sets.

It is well known that the classical theory of fuzzy sets is a basis for the theory of possibility [30, 131]. Of course, it is possible to develop a similar generalized theory of possibility starting from generalized fuzzy sets, see, e.g., [30, 59, 69]. Generalized theories can be noncommutative; they seem to be more qualitative and less quantitative with respect to the classical theories presented in [130, 131]. We see that idempotent analysis and the theories of (generalized) fuzzy sets and possibility have the same objects, i.e. functions taking their values in semirings. However, basic problems and methods could be different for these theories (like for the measure theory and the probability theory).

14. Relations to other areas and miscellaneous applications

Many relations and applications of idempotent mathematics to different theoretical and applied areas of mathematical sciences are discussed above. Of course, optimization and optimal control problems form a very natural field for applications of ideas and methods of idempotent mathematics. There is a very good survey paper [62] of V. N. Kolokoltsov on the subject, see also [6, 11, 13, 16, 19–22, 24, 26, 27, 34–37, 46–48, 70, 72, 82, 83, 85, 87–92, 106, 125–127, 129, 136, 137].

There are many applications to differential equations and stochastic differential equations, see, e.g., [34–37, 48, 60, 61, 63, 65, 87–91, 100, 118, 119].

Applications to the game theory are discussed, e.g., in [64, 65, 90]. There are interesting applications in biology (bioinformatics), see, e.g., [33, 99, 112]. Applications and relations to mathematical morphology are examined the paper [27] of P. Del Moral and M. Doisy and especially in an extended preprint version of this article. There are many relations and applications to physics (quantum and classical physics, statistical physics, cosmology etc.) see, e.g., [15, 61, 65, 78, 79, 98, 107], section 12 above and the paper of P. Lotito, J.-P. Quadrat, E. Mancinelli [85] published in this volume.

There are important relations and applications to purely mathematical areas. The so-called tropical combinatorics is discussed in a large survey paper [57] of A. N. Kirillov, see also [11, 137]. Tropical mathematics is closely related to the very attractive and popular theory of cluster algebras founded by S. Fomin and A. Zelevinsky, see their survey paper [39]. In both cases there are relations with the traditional theory of representations of Lie groups and related topics. There are important relations with convex analysis and discrete convex analysis, see, e.g., [2, 18, 21, 28, 81, 86, 91, 116, 133–135] and the paper of G. Cohen, S. Gaubert, J.-P. Quadrat, and J. Singer published in the present volume.

Many authors examine, explicitly or not, relations and applications of idempotent mathematics to mathematical economics starting from the classical papers of N. N. Vorobjev [125–127], see, e.g., [23, 64, 91, 132, 137].

References

[1] M. Akian, *Densities of idempotent measures and large deviations*, Trans. Amer. Math. Soc. **351** (1999), 4515–4543.

[2] M. Akian and S. Gaubert, *Spectral theorem for convex monotone homogeneous maps and ergodic control*, Nonlinear Analysis **52** (2003), 637–679. See also arXiv:math.SP/0110108.

[3] M. Akian, S. Gaubert, and V. Kolokoltsov, *Set Coverings and Invertibility of Functional Galois Connections.* – In [73], this volume.

[4] M. Akian, J. P. Quadrat, and M. Viot, *Duality between probability and optimization.* — In [48], p. 331–353.

[5] M. Akian, S. Gaubert, and C. Walsh, *Discrete Max-Plus Spectral Theory.* – In [73], this volume.

[6] F. Baccelli, G. Cohen, G. J. Olsder, and J.-P. Quadrat, *Synchronization and Linearity: An Algebra for Discrete Event Systems*, John Wiley & Sons Publishers, New York e.a., 1992.

[7] M. Bardi, I. Capuzzo-Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton–Jacobi–Bellman Equations*, Birkhäuser, Boston–Basel–Berlin, 1997.

[8] P. Bernhard, *Minimax versus stochastic partial information control*, Proceedings of the 33rd Conference on Decision and Control, Lake Buena Vista, FL, December 1994, IEEE, 1994, p. 2572–2577.

[9] Y. Brenier, U. Frisch, M. Hénon, G. Loeper, S. Matarrese, R. Mohayaee, and A. Sobolevskiĭ. *Reconstruction of the early Universe as a convex optimization problem*, Mon. Nat. R. Astron. Soc. **346** (2003), 501–524.

[10] P. Butkovič, *Strong regularity of matrices — a survey of results*, Discrete Applied Math. **48** (1994), 45–68.

[11] P. Butkovič, *On the combinatorial aspects of max-algebra.* – In [73], this volume.

[12] I. Capuzzo Dolcetta, P.-L. Lions (Eds.), *Viscosity solutions and applications*, Lectures given at the 2nd Session of the C.I.M.E. held in Montecatini Terme, Italy, June 12–20, 1995. Lecture Notes in Mathematics **1660**, 1997.

[13] B. A. Carré, *Graphs and networks*, The Clarendon Press/Oxford University Press, Oxford, 1979.

[14] K. Cechlárová and R. A. Cuninghame-Green, *Interval systems of max-separable linear equations*, Linear Algebra and its Applications **340** (2002), 215–224.

[15] Weiren Chou and R. J. Duffin, *An additive eigenvalue problem of physics related HJB equation to linear programming*, Adv. in Applied Mathematics **8** (1987), 486–498.

[16] G. Cohen, S. Gaubert, and J. P. Quadrat, *Max-plus algebra and system theory: where we are and where to go now*, Annual Reviews in Control **23** (1999), 207–219.

[17] G. Cohen, S. Gaubert, and J.-P. Quadrat, *Duality and separation theorems in idempotent semimodules*, Linear Algebra and its Applications **379** (2004), 395–422. Also arXiv:math.FA/0212294.

[18] G. Cohen, S. Gaubert, J.-P. Quadrat, and I. Singer, *Max-plus convex sets and functions.* — In [73], this volume.

[19] G. Cohen and J.-P. Quadrat (Eds.), *11th International Conference on Analysis and Optimization Systems*, Springer Lecture Notes on Control and Information Systems **199**, 1994.

[20] R. A. Cuninghame-Green, *Minimax algebra*, Springer Lect. Notes in Economics and Mathematical Systems **166**, Berlin et al., 1979.

[21] R. A. Cuninghame-Green, *Minimax algebra and applications*, Advances in Imaging and Electron Physics **90** (1995), 1–121. (Academic Press, New York).

[22] R. Cuninghame-Green and P. Meijer, *An algebra for piesewise-linear minimax problems*, Dicsrete Appl. Math. **2** (1980), 267–294.

[23] V. Danilov, G. Koshevoi, and K. Murota, *Discrete convexity and equilibria in economics with indivisible goods and money*, Math. Soc. Sci. **11** (2001), 251–273.

[24] P. Del Moral, *A survey of Maslov optimization theory.* – In: V. N. Kolokoltsov and V. P. Maslov, *Idempotent Analysis and Applications*, Kluwer Acad. Publ., Dordrecht, 1997, p. 243–302 (Appendix).

[25] P. Del Moral, *Feynman–Kac formulae. Genealogical and interacting particle systems with applications*, Springer, New York e.a., 2004.

[26] P. Del Moral and M. Doisy, *Maslov idempotent probability calculus, I, II*, Theory of Probability and its Applications **43** (1998), no. 4, 735–751 and **44** (1999), no. 2, 384–400.

[27] P. Del Moral and M. Doisy, *On the applications of Maslov optimization theory*, Mathematical Notes **69** (2001), no. 2, 232–244.

[28] M. Develin and B. Sturmfels, *Tropical covexity*, arXiv:math.MG/0308254, 2003.

[29] A. Di Nola, B. Gerla, *Algebras of Lukasiewicz's Logic and Their Semiring Reducts..* – In [73], this volume.

[30] D. Dubois, H. Prade, and R. Sabbadin, *Decision-theory foundations of qualitative possibility theory*, European Journal of Operational Research **128** (2001), 459–478.

[31] P. S. Dudnikov and S. N. Samborskiĭ, *Endomorphisms of semimodules over semirings with an idempotent operation*, preprint of the Mathematical Institute of the Ukrainian Academy of Sciences, Kiev, 1987 (in Russian); Izv. Akad. Nauk SSSR, ser. math. **55** (1991), no. 1, 93–109; English transl. in Math. USSR Izvestiya **38** (1992), no. 1, 91–105.

[32] M. Einsiedler, M. Kapranov, and D. Lind, *Non-archimedean amoebas and tropical varieties*, arXiv:math.AG/0408311, 2004.

[33] A. V. Finkelstein and M. A. Roytberg, *Computation of biopolymers: a general approach to different problems*, BioSystems **30** (1993), 1–20.

[34] W. H. Fleming, *Max-Plus Stochastic Control.* — In: B. Pasik-Duncan, Ed., *Stochastic Theory and Control*, Lecture Notes in Control and Information Science **280** (2002), 111–119.

[35] W. H. Fleming, *Max-plus stochastic processes*, Applied Math. And Optim. **48** (2004), 159–181.

[36] W. H. Fleming and W. M. McEneaney, *A max-plus-based algorithm for a Hamilton–Jacobi–Bellman equation of nonlinear filtering*, SIAM J. Control Optim. **38** (2000), no. 3, 683–710.

[37] W. H. Fleming and W. M. McEneaney, *Max-plus approaches to continuous space control and dynamic programming.* — In [73], this volume.

[38] W. H. Fleming and H. M. Soner, *Controlled Markov processes and viscosity solutions*, Springer, New York, 1993.

[39] S. Fomin and A. Zelevinsky, *Cluster algebras: Notes for the CDM-03 Conference*, Conf. "Current Developments in Mathematics 2003" held at Harvard University on November 21–22, 2004, arXiv:math.RT/0311493, v. 2, 2004.

[40] U. Frisch, S. Matarrese, R. Mohayaee, and A. Sobolevskiĭ. *A reconstruction of the initial conditions of the Universe by optimal mass transportation*, Nature, **417** (May 2002), 260–262.

[41] I. M. Gelfand, M. M. Kapranov, and A. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Birkhäuser, Boston, 1994.

[42] K. Glazek, *A guide to the literature on semirings and their applications in mathematics and information sciences: with complete bibliography*, Kluwer Acad. Publ., Dordrecht e.a., 2002.

[43] J. A. Goguen, *L-fuzzy sets*, J. of Math. Anal. Appl. **18** (1967), no. 1, 145–174.

[44] J. S. Golan, *Semirings and their applications*, Kluwer Acad. Publ., Dordrecht, 1999.

[45] J. S. Golan, *Semirings and affine equations over them: thery and applications*, Kluwer Acad. Publ., Dordrecht, 2003.

[46] M. Gondran and M. Minoux, *Graphes et algorithmes*, Editions Eyrolles, Paris, 1979, 1988.

[47] M. Gondran and M. Minoux, *Graphes, dioïdes et semi-anneaux*, Editions TEC&DOC, Paris e.a., 2001.

[48] J. Gunawardena (Ed.), *Idempotency*, Publ. of the Newton Institute, Vol. **11**, Cambridge University Press, Cambridge, 1998.

[49] J. Gunawardena, *An introduction to idempotency.* — In [48], p. 1–49.

[50] I. Itenberg, V. Kharlamov, and E. Shustin, *Welschinger invariant and enumeration of real rational curves*, International Mathematics Research Notices **49** (2003), 26–39.

[51] I. Itenberg, V. Kharlamov, and E. Shustin, *Logarithmic equivalence of Welschinger and Gromov–Witten invariants*, arXiv:math.AG/0407188, 2004.

[52] M. M. Kapranov, *Amoebas over non-Archimedian fields*, Preprint, 2000.

[53] Y. Katsov, *On flat semimodules over semirings*, Algebra Universalis **51** (2004), 287–299.

[54] K. Khanin, D. Khmelëv, and A. Sobolevskiĭ, *A blow-up phenomenon in the Hamilton–Jacobi equation in an unbounded domain* – In [73], this volume.

[55] K. Khanin, D. Khmelëv, and A. Sobolevskiĭ, *On velocities of lagrangian minimizers*. Moscow Math. J., 2005 (in print).

[56] K. H. Kim and F. W. Roush, *Inclines and incline matrices: a survey*, Linear Algebra and its Applications **379** (2004), 457–473.

[57] A. N. Kirillov, *Introduction to tropical combinatorics*. – In: A. N. Kirillov and N. Liskova (Eds.), *Physics and Combinatorics 2000*, Proc. of the Nagoya 2000 Intern. Workshop, World Scientific, 2001, p. 82–150.

[58] S. C. Kleene, *Representation of events in nerve sets and finite automata*. – In: J. McCarthy and C. Shannon (Eds), Automata Studies, Princeton University Press, Princeton, 1956, pp. 3–40.

[59] E. P. Klement and E. Pap (Eds.), *Mathematics of Fuzzy Systems*, 25th Linz Seminar on Fuzzy Set Theory, Linz, Austria, Feb. 3–7, 2004. Abstracts. J. Kepler Univ., Linz, 2004.

[60] V. N. Kolokoltsov, *Stochastic Hamilton–Jacobi–Bellman equations and stochastic Hamiltonian systems*, Journal of Control and Dynamic Systems **2** (1996), no. 3, 299–319.

[61] V. N. Kolokoltsov, *Semiclassical analysis for diffusions and stochastic processes*, Springer Lecture Notes in Math. **1724**, 2000.

[62] V. N. Kolokoltsov, *Idempotency structures in optimization*, Journal Math. Sci. **104** (2001), no. 1, 847–880.

[63] V. N. Kolokoltsov, *Small diffusion and fast dying out asymptotics for superprocesses as non-Hamiltonian quasiclassics for evalution equations*, Electronic Journal of Probability **6** (2001), paper 21 (see http://www.math.washington.edu/∼ ejpecp/).

[64] V. N. Kolokoltsov and O. A. Malafeyev, *A turnpike theorem in conflict-control processes with many participants*. — In: O. Malafeyev (Ed.), *Conflict Models in Economics and Finance*, St. Petersburg Univ. Press, 1997 (in Russian).

[65] V. Kolokoltsov and V. Maslov, *Idempotent analysis and applications*, Kluwer Acad. Publ., 1997.

[66] V. Kolokoltsov and A. Tyukov, *Small time amd semiclassical asymptotics for stochastic heat equation driven by Levi noise*, Stochastics and Stochastics Reports **75** (2003), no. 1–2, 1–38.

[67] M. Kontsevich and Y. Soibelman, *Homological mirror symmetry and torus fibration.* — In: Symplectic geometry and mirror symmetry (Seoul, 2000), World Sci. Publ., River Edge, N.J., 2001, pp. 203–263.

[68] M. Kontsevich and Y. Soibelman, *Affine structures and non-archimedean analytic spaces*, arXiv:math.AG/0406564, 2004.

[69] G. L. Litvinov, *Dequantization of mathematics, idempotent semirings and fuzzy sets.* – In [59], p. 113–117.

[70] G. L. Litvinov and V. P. Maslov, *Correspondence principle for idempotent calculus and some computer applications*, (IHES/M/95/33), Institut des Hautes Etudes Scientifiques, Bures-sur-Yvette, 1995. Also arXiv:math.GM/0101021.

[71] G. L. Litvinov and V. P. Maslov, *Idempotent mathematics: correspondence principle and applications*, Russian Mathematical Surveys **51** (1996), no. 6, 1210–1211.

[72] G. L. Litvinov and V. P. Maslov, *The correspondence principle for idempotent calculus and some computer applications.* — In [48], p. 420–443.

[73] G. L. Litvinov and V. P. Maslov (Eds.), *Idempotent mathematics and mathematical physics*, Contemporary Mathematics, AMS, Providence, RI, 2005 (this volume).

[74] G. L. Litvinov and E. V. Maslova, *Universal numerical algorithms and their software implementation*, Programming and Computer Software **26** (2000), no. 5, 275–280. Also arXiv:math.SC/0102114.

[75] G. L. Litvinov, V. P. Maslov, and A. Ya. Rodionov, *A unifying approach to software and hardware design for scientific calculations and idempotent mathematics*, International Sophus Lie Centre, Moscow 2000. Also arXiv:math.SC/0101069.

[76] G. L. Litvinov, V. P. Maslov, and G. B. Shpiz, *Linear functionals on idempotent spaces: an algebraic approach*, Doklady Mathematics **58** (1998), no. 3, 389–391. Also arXiv:math.FA/0012268.

[77] G. L. Litvinov, V. P. Maslov, and G. B. Shpiz, *Tensor products of idempotent semimodules. An algebraic approach*, Mathematical Notes **65** (1999), no. 4, 497–489. Also arXiv:math.FA/0101153.

[78] G. L. Litvinov, V. P. Maslov, and G. B. Shpiz, *Idempotent functional analysis. An algebraic approach*, Mathematical Notes **69** (2001), no. 5, 696–729. Also arXiv:math.FA/0009128.

[79] G. L. Litvinov, V. P. Maslov, and G. B. Shpiz, *Idempotent (asymptotic) analysis and the representation theory.* – In: V. A. Malyshev and A. M. Vershik (Eds.), Asymptotic Combinatorics with Applications to Mathematical Physics. Kluwer Academic Publ., Dordrecht et al, 2002, p. 267–278. Also arXiv:math.RT/0206025.

[80] G. L. Litvinov and G. B. Shpiz, *Nuclear semimodules and kernel theorems in idempotent analysis: an algebraic approach*, Doklady Mathematics **66** (2002), no. 2, 197–199. Also arXiv :math.FA/0202026.

[81] G. L. Litvinov and G. B. Shpiz, *Dequantization transform and generalized Newton polytopes.* — In [73], this volume.

[82] G. L. Litvinov and A. N. Sobolevskiĭ, *Exact interval solutions of the discrete Bellman equation and polynomial complexity of problems in interval idempotent linear algebra*, Doklady Mathematics **62** (2000), no. 2, 199–201. Also arXiv:math.LA/0101041.

[83] G. L. Litvinov and A. N. Sobolevskiĭ, *Idempotent interval analysis and optimization problems*, Reliable Computing **7** (2001), no. 5, 353–377. Also arXiv:math.SC/0101080.

[84] P. Loreti and M. Pedicini, *An object-oriented approach to idempotent analysis: integral equations as optimal control problems.* — In [73], this volume.

[85] P. Lotito, J.-P. Quadrat, and E. Mancinelli, *Traffic assignment* & *Gibbs–Maslov semirings.* — In [73], this volume.

[86] G. G. Magaril-Il'yaev and V. M. Tikhomirov, *Convex analysis: theory and applications*, Translations of Mathematical Monographs, vol. **222**, American Math. Soc., Providence, RI, 2003.

[87] V. P. Maslov, *New superposition principle for optimization problems.* — In: Seminaire sur les Equations aux Dérivées Partielles 1985/86, Centre Math. De l'Ecole Polytechnique, Palaiseau, 1986, exposé 24.

[88] V. P. Maslov, *On a new superposition principle for optimization problems*, Uspekhi Mat. Nauk, [Russian Math. Surveys], **42**, no. 3 (1987), 39–48.

[89] V. P. Maslov, *Méthodes opératorielles*, Mir, Moscow, 1987.

[90] V. P. Maslov and V. N. Kolokoltsov, *Idempotent analysis and its application in optimal control*, Nauka, Moscow, 1994. (in Russian)

[91] V. P. Maslov and S. N. Samborskiĭ (Eds), *Idempotent analysis*, Adv. in Sov. Math., vol. **13**, AMS, RI, 1992.

[92] V. P. Maslov and K. A. Volosov (Eds.), *Mathematical aspects of computer engineering*, MIR Publ., Moscow, 1988.

[93] D. McCaffrey, *Viscosity solutions on Lagrangian manifolds and connections with tunneling operators.* — In [73], this volume.

[94] G. Mikhalkin, *Amoebas of algebraic varieties*, Notes for the Real Algebraic and Analytic Geometry Congress, June 11–15, 2001, Rennes, France. Also arXiv:math.AG/0108225, 2001.

[95] G. Mikhalkin, *Counting curves via lattice path in polygons*, C.R. Acad. Sci. Paris **336** (2003), no. 8, 629–634.

[96] G. Mikhalkin, *Amoebas of algebraic varieties and tropical geometry*, to be published in volume "Different faces in Geometry." Also arXiv:math.AG/0403015, 2004.

[97] G. Mikhalkin, *Enumerative tropical algebraic geometry in* $\mathbf{R}^2$, Journal of the ACM, 2005 (in press). Also arXiv:math.AG/0312530.

[98] R. D. Nussbaum, *Convergence of iterates of a nonlinear operator arising in statistical mechanics*, Nonlinearity **4** (1991), 1223–1240.

[99] L. Pachter and B. Sturmfels, *The mathematics of phylogenomics*, arXiv:math.ST/0409132, 2004.

[100] E. Pap, *Applications of the generated pseudo-analysis to nonlinear partial differential equations.* – In [73], this volume.

[101] M. Passare and A. Tsikh, *Amoebas: their spines and their contours.* — In [73], this volume.

[102] J. E. Pin, *Tropical semirings.* – In [48], p. 50–60.

[103] A. A. Puhalskii, *Large Deviations and Idempotent Probability*, Chapman and Hall/CRC Press, London/Boka Raton, FL., 2001.

[104] A. A. Puhalskii, *On large deviations convergence of invariant measures*, Journal of Theoretical Probability **16** (2003), 689–724.

[105] J.-P. Quadrat, *Théorèmes asymptotiques en programmation dynamique*, Comptes Rendus Acad. Sci., Paris **311**, (1990), 745–748.

[106] J.-P. Quadrat and Max-Plus working group, *Max-plus algebra and applications to system theory and optimal control*, – In: *Proc. of the Internat. Congress of Mathematicians*, Zürich, 1994.

[107] J.-P. Quadrat and Max-Plus working group, *Min-plus linearity and statistical mechanics*, Markov Processes and Related Fields **3** (1997), no. 4, 565–587.

[108] J. Richter-Gebert, B. Sturmfels, and T. Theobald, *First steps in tropical geometry.* — In [73], this volume. Also arXiv:math.AG/0306366.

[109] K. I. Rosenthal, *Quantales and their applications*, Pitman Research Notes in Mathematics Series, Longman Sci.&Tech., 1990.

[110] K. I. Rosenthal, *The theory of quantaloids*, Pitman Research Notes in Mathematics Series **348**, LongmanHarlow, 1996.

[111] I. V. Roublev, *On minimax and idempotent generalized weak solutions to the Hamilton–Jacobi Equation.* – In [73], this volume.

[112] M. A. Roytberg, *Fast algorithm for optimal aligning of symbol sequences*, DIMACS Series in Mathematics and Theoretical Computer Science, **8** (1992), 113–126.

[113] H. Rullgård, *Polynomial amoebas and convexity*, Preprint, Stokholm University, 2001.

[114] G. B. Shpiz, *Solution of algebraic equations over idempotent semifields*, Uspekhi Matem. Nauk [Russian Math. Surveys] **55** (2000), no. 5, 185–186.

[115] E. Shustin, *Patchworking singular algebraic curves, non Archimedian amoebas and enumerative geometry*, arXiv:math.AG/0211278.

[116] I. Singer, *Abstract convex analysis*, John Wiley&Sons Inc., New York, 1997.

[117] I. Singer, *Some relations between linear mappings and conjugations in idempotent analysis*, J. Math. Sci. **115** (2003), no. 5, 2610–2630.

[118] A. N. Sobolevskiĭ, *Aubry-Mather theory and idempotent eigenfunctions of Bellman operator*, Commun. Contemp. Math. **1** (1999), no. 4, 517–533.

[119] A. N. Sobolevskiĭ, *Periodic solutions of the Hamilton-Jacobi equation with a periodic nonhomogeneity, and the Aubry-Mather theory*, Mat. Sb. **190** (1999), no. 10, 87–104.

[120] D. Speyer and B. Sturmfels, *Tropical mathematics*, arXiv:math.CO/0408099, 2004.

[121] B. Sturmfels, *Solving systems of polynomial equations*, CBMS Regional Conference Series in Mathematics, AMS, Providence, RI, 2002.

[122] A. I. Subbotin, *Generalized solutions of first order PDEs: The dynamic optimization perspective*. Birkhäuser, Boston, 1996.

[123] O. Viro, *Dequantization of real algebraic geometry on a logarithmic paper*. — In: 3rd European Congress of Mathematics, Barcelona, 2000. Also arXiv:math/0005163.

[124] O. Viro, *What is an amoeba?*, Notices of the Amer. Math, Soc. **49** (2002), 916–917.

[125] N. N. Vorobjev, *The extremal matrix algebra*, Soviet Math. Dokl. **4** (1963), 1220–1223.

[126] N. N. Vorobjev, *Extremal algebra of positive matrices*, Elektronische Informationsverarbeitung und Kybernetik **3** (1967), 39–57. (in Russian)

[127] N. N. Vorobjev, *Extremal algebra of nonnegative matrices*, Elektronische Informationsverarbeitung und Kybernetik **6** (1970), 302–312. (in Russian)

[128] E. Wagneur, *Dequantization: semi-direct sums of idempotent semimodules*. - In [73], this volume.

[129] J. van der Woude and G. J. Olsder, *On* $(\min, \max, +)$-*inequalities*.-In [73], this volume.

[130] L. A. Zadeh, *Fuzzy sets*, Information and Control **8** (1965), 338–353.

[131] L. A. Zadeh, *Fuzzy sets as a basis for a theory of possibility*, Fuzzy Sets and Systems, **1** (1978), 3–28.

[132] K. Zimmermann, *Extremální Algebra*, Ekonomický ùstav ČSAV, Praha **46** (1976). (in Czech).

[133] K. Zimmermann, *A general separation theorem in extremal algebras*, Ekonom.-Mat. Obzor **13** (1977), no. 2, 179–210.

[134] K. Zimmermann, *Extremally convex functions*, Wiss. Z. Päd. Hochschule "N. K. Krupskaya" **17** (1979), 147–158.

[135] K. Zimmermann, *A generalization of convex functions*, Ekonom.-Mat. Obzor **15** (1979), no. 2, 147–158.

[136] K. Zimmermann, *Solution of some max-separable optimization problems with inequality constraints.* — In [73], this volume.

[137] U. Zimmermann, *Linear and combinatorial optimization in ordered algebraic structures*, Ann. Discrete Math., **10** (1981), 1–380.

INDEPENDENT UNIVERSITY OF MOSCOW, BOL'SHOI VLASIEVSKII PER., 11, MOSCOW 121002, RUSSIA

E-mail address: `islc@dol.ru`